

## **PREGALIN X 75**

### **For the Use of a Registered Medical Practitioner or a Hospital or a Laboratory Only**

Abbreviated Prescribing information for Pregalin X (Pregabalin 75mg & Methylcobalamin 750mcg Capsules) [Please refer the complete prescribing information available at [www.torrentpharma.com](http://www.torrentpharma.com)]

**PHARMACOLOGICAL PROPERTIES:** *Pregabalin*: It is a structural derivative of the inhibitory neurotransmitter gammaaminobutyric acid (GABA). *Methylcobalamin* is required for synthesis of deoxyribonucleic acid.

**INDICATION:** It is indicated for the treatment of peripheral neuropathy in adult patients.

**DOSAGE AND ADMINISTRATION:** It should be taken twice daily.

**CONTRAINDICATION:** It is contraindicated in patients with known hypersensitivity to any of its components.

**WARNINGS & PRECAUTIONS:** Caution should be taken while withdrawal OR rapid discontinuation of the treatment. It may cause dizziness, somnolence, ophthalmological effects or reactions, weight gain, peripheral edema, creatine kinase elevations, decrease in platelets count, PR interval prolongation and ECG changes.

**ADVERSE REACTIONS:** Dizziness, somnolence, asthenia, confusion, peripheral edema, ataxia, abnormal gait, infection, flu syndrome, dry mouth, myasthenia, bronchitis, diplopia, urinary incontinence, allergic reaction, suicide attempt, cellulitis, pelvic pain, deep thrombophlebitis, heart failure, gastroenteritis, gastrointestinal hemorrhage, increased appetite, glucose tolerance decreased, ecchymosis, anxiety, libido decreased, apnea, stevens-johnson syndrome, lung fibrosis, anorgasmia, impotence, nausea, vomiting and diarrhea.

### **MARKETED BY:**



**TORRENT PHARMACEUTICALS LTD.**

Torrent House, Off Ashram Road,

Ahmedabad-380 009, INDIA

**IN/ PREGALIN X/75mg,7500mcg/MAR 2015/01/AbPI**

(Additional information is available on request)