

PREGEB OD

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory Only

Abbreviated Prescribing information for **Pregeb OD** [Pregabalin Sustained Release Tablets 75,150 and 300mg] [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Pregabalin binds with high affinity to the alpha2-delta site (an auxiliary subunit of voltage gated calcium channels) in central nervous system tissues and it is a structural derivative of the inhibitory neurotransmitter gamma amino butyric acid (GABA) and thus produces anti-seizure effect and relief against neuropathic pain.

INDICATION: Pregabalin Sustained Release tablet is indicated for management of neuropathic pain.

DOSAGE AND ADMINISTRATION: The starting dose of Pregabalin SR Tablets is 150 mg once daily, administered in the evening. Maximum safe dose of Pregabalin is 600 mg/day. Pregabalin SR Tablets can be taken with or without food. Pregabalin SR must be swallowed whole with the aid of liquids. Tablets should not be chewed, divided, or crushed.

CONTRAINDICATION: Contraindicated in patients with known hypersensitivity to pregabalin or any components of this product.

WARNINGS & PRECAUTIONS: Angioedema, hypersensitivity, Pregabalin should be withdrawn gradually to minimize the potential of increased seizure frequency in patients with seizure disorders and in some patients reported symptoms including insomnia, nausea, headache and diarrhea. Peripheral Edema dizziness and somnolence, blurred vision, weight gain, peripheral edema, creatine kinase elevations, decreased platelet count and PR interval prolongation.

DRUG INTERACTIONS: No specific drug interactions.

ADVERSE REACTIONS: Dizziness, somnolence, dry mouth, edema, blurred vision, weight gain, thinking abnormal, asthenia, back pain, chest pain, constipation, gastroenteritis, increased appetite, flatulence, peripheral edema, weight gain, hypoglycemia, neuropathy, ataxia, vertigo, confusion, euphoria, incoordination, abdominal pain, tremor, abnormal gait, amnesia, accidental injury, nervousness, dyspnea, abnormal vision, abdominal pain, allergic reaction, fever, gastroenteritis, increased appetite, ecchymosis, arthralgia, leg cramps, myalgia, myasthenia, anxiety, depersonalization, deep thrombophlebitis, heart failure, hypotension, postural hypotension, hypertonia, hypoesthesia, libido decreased, nystagmus, paresthesia, sedation, stupor, twitching; abnormal dreams, agitation, apathy, aphasia, hallucinations, conjunctivitis, diplopia, anorgasmia, impotence, urinary frequency, urinary incontinence, pulmonary edema, congestive heart failure, diarrhea itching; transitory exanthema, blood clots, itching and serious allergic reactions.

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Torrent House, Off Ashram Road,
Ahmedabad-380 009, INDIA

IN/ PREGEB OD 75,150,300mg/Aug-15/01/ABPI

(Additional information is available on request)