

DROXYL 250 INSTA USE 30 ML

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory Only

Abbreviated Prescribing information for DROXYL 250 INSTA USE 30 ML [Cefadroxil 250 mg/5ml]
[Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Beta-lactam antibiotics, cephalosporins which inhibits bacterial wall synthesis of actively dividing cells by binding to one or more penicillin-binding proteins.

INDICATION: Indicated in the treatment of urinary tract infections, skin and skin structure infection caused by *S. Cocci*, pharyngitis tonsillitis, respiratory tract infections only.

DOSAGE AND ADMINISTRATION: As per the physician direction.

CONTRAINDICATION: Hypersensitivity to any of the components of product or to cephalosporin group antibiotics.

WARNINGS & PRECAUTIONS: Previous hypersensitivity to cephalosporin, or penicillins; creatinine clearance less than 50ml/min; Prolong use may result in overgrowth of nonsusceptible organism; Colitis; False positive laboratory tests (i.e. positive comb's test) reported; use during pregnancy and to lactating mother's.

DRUG INTERACTIONS: It is having interaction with bacteriostatic antibiotics like tetracycline, erythromycin, sulfonamides, chloramphenicol; also nephrotoxic agents like aminoglycoside antibiotics, polymyxin B, colistin or high-dose loop diuretics; anticoagulants or thrombocyte aggregation inhibitors (long term use); probenecid; oral contraceptives; cholestyramine; false positive glycosuria tests (with Benedict's or Fehling's solutions).

ADVERSE REACTIONS: Hyper sensitivity reactions (Pruritus, rash, allergic exanthema, urticarial, angioneurotic oedema, drug fever, serum sickness-like reactions, arthralgia, interstitial nephritis, Immediate allergic reaction, SJS, erythema multiforme), eosinophilia, thrombocytopenia, leucopenia, neutropenia, agranulocytosis, haemolytic anaemia, nausea, vomiting, diarrhoea, dyspepsia, abdominal pain, glossitis, pseudomembranous colitis, *C. Difficile* diarrhea, elevation of serum transaminases, ALP, cholestase, idiosyncratic hepatic failure, headache, dizziness, nervousness, sleeplessness, fatigue, vaginal mycoses, thrush, colitis.

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Torrent House, Off Ashram Road,
Ahmedabad-380 009, INDIA

IN/DROXYL 250 INSTA USE 30 ML/Mar-2015/01/AbPI

(Additional information is available on request)