

**For the use of a Registered Medical Practitioner or Hospital or a Laboratory only**

---

**TRIVOGLITOR FORTE**

(Metformin Hydrochloride (SR), Glimepiride and Voglibose Tablets)

---

**COMPOSITION**

**TRIVOGLITOR FORTE 1**

Each uncoated bilayered tablet contains:

Metformin Hydrochloride I. P. 500 mg (in sustained release form)

Glimepiride I.P. 1 mg

Voglibose I.P. 0.3 mg

Color: Sunset yellow Lake

**TRIVOGLITOR FORTE 2**

Each uncoated bilayered tablet contains:

Metformin Hydrochloride I.P. 500 mg (in sustained release form)

Glimepiride IP. 2 mg

Voglibose I.P. 0.3 mg

Color: Quinoline Yellow Lake

**INDICATIONS**

Trivogliton Forte is indicated in the treatment of patients with type II diabetes mellitus when diet, exercise and therapy with two agents do not result in adequate glycaemic control.

**POSODOLOGY AND METHOD OF ADMINISTRATION**

Trivogliton Forte should be given once daily before a full meal

Therapy should be started with Trivogliton Forte 1 or 2 one tablet daily based on previous drug treatment and glycaemic status of the patient. Depending upon the glycaemic control, dose may be titrated to 2 tablets once daily.

The basis for successful treatment of diabetes is a good diet, regular physical activity, as well as routine checks of blood and urine. Tablets or insulin cannot compensate if the patient does not keep to the recommended diet. Dosage is determined by the results of blood and urinary glucose determinations.

**CONTRAINDICATIONS**

Trivogliton contraindicated in the following patients

1. Patients with a history of hypersensitivity to Metformin hydrochloride, Glimepiride and Voglibose or any of the ingredients of this product.
2. Renal disease or renal dysfunction (e.g., as suggested by serum creatinine levels 1.5 mg/ dL [males], 1.4 mg/dL [females] or abnormal creatinine clearance), which may also result from conditions such as cardiovascular collapse (shock), acute myocardial infarction, and septicemia.
3. Congestive heart failure requiring pharmacological treatment.
4. Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma. Diabetic ketoacidosis should be treated with insulin.

5. Sulfonamide derivatives: Patients who have developed an allergic reaction to sulfonamide derivatives may develop an allergic reaction to Glimepiride. Do not use Glimepiride in patients who have a history of an allergic reaction to sulfonamide derivatives. Reported hypersensitivity reactions include cutaneous eruptions with or without pruritus as well as more serious reactions (e.g. anaphylaxis, angioedema, Stevens-Johnson syndrome, and dyspnea)

6. Patients with severe ketosis, or in a state of diabetic coma or pre-coma (since it becomes essential to quickly rectify hyperglycaemia with administration of intravenous fluid or insulin, the use of voglibose is not suitable).

7. Patients with severe infections, before and after operation, or with serious trauma (It is desirable to control blood sugar with the insulin. Therefore, administration of the drug is not appropriate).

## **SPECIAL WARNINGS AND PRECAUTIONS FOR USE**

### **Glimepiride**

It must be taken shortly before or during a meal.

When meals are taken at irregular hours or skipped altogether, treatment with Glimepiride may lead to hypoglycaemia. Possible symptoms of hypoglycaemia include: headache, ravenous hunger, nausea, vomiting, lassitude, sleepiness, disordered sleep, restlessness, aggressiveness, impaired concentration, alertness and reaction time, depression, confusion, speech and visual disorders, aphasia, tremor, paresis, sensory disturbances, dizziness, helplessness, loss of self-control, delirium, cerebral convulsions, somnolence and loss of consciousness up to and including coma, shallow respiration and bradycardia. In addition, signs of adrenergic counter-regulation may be present such as sweating, clammy skin, anxiety, tachycardia, hypertension, palpitations, angina pectoris and cardiac arrhythmias. The clinical picture of a severe hypoglycaemic attack may resemble that of a stroke.

Symptoms can almost always be promptly controlled by immediate intake carbohydrates (sugar). Artificial sweeteners have no effect. It is known from other sulfonylureas that, despite initially successful countermeasures, hypoglycaemia may recur. Severe hypoglycaemia or prolonged hypoglycaemia, only temporarily controlled by the usual amounts of sugar, require immediate medical treatment and occasionally hospitalisation.

Factors favouring hypoglycaemia include:

- unwillingness or (more commonly in older patients) incapacity of the patient to cooperate,
- undernutrition, irregular mealtimes or missed meals or periods of fasting,
- alterations in diet,
- imbalance between physical exertion and carbohydrate intake,
- consumption of alcohol, especially in combination with skipped meals,
- impaired renal function,
- serious liver dysfunction,
- overdose with Glimepiride,
- certain uncompensated disorders of the endocrine system affecting carbohydrate metabolism or counter-regulation of hypoglycaemia (as for example in certain disorders of thyroid function and in anterior pituitary or adrenocortical insufficiency),
- concurrent administration of certain other medicinal products

Treatment with Glimpiride requires regular monitoring of glucose levels in blood and urine. In addition determination of the proportion of glycosylated haemoglobin is recommended.

Regular hepatic and haematological monitoring (especially leucocytes and thrombocytes) are required during treatment with Glimpiride.

In stress-situations (e.g. accidents, acute operations, infections with fever, etc.) a temporary switch to insulin may be indicated.

No experience has been gained concerning the use of Glimpiride in patients with severe impairment of liver function or dialysis patients. In patients with severe impairment of renal or liver function change over to insulin is indicated.

Treatment of patients with G6PD-deficiency with sulfonylurea agents can lead to hemolytic anaemia. Since glimepiride belongs to the class of sulfonylurea agents, caution should be used in patients with G6PD-deficiency and a non-sulfonylurea alternative should be considered.

### **Metformin**

Lactic acidosis:

Lactic acidosis is a very rare, but serious (high mortality rate in the absence of prompt treatment), metabolic complication that can occur due to metformin accumulation. Reported cases of lactic acidosis in patients on metformin have occurred primarily in diabetic patients with impaired renal failure or acute worsening of renal function. Special caution should be paid to situations where renal function may become impaired, for example in case of dehydration (severe diarrhea or vomiting), or when initiating antihypertensive therapy or diuretic therapy and when starting therapy with a non-steroidal anti-inflammatory drug (NSAID). In the acute conditions listed, metformin should be temporarily discontinued.

Other associated risk factors should be considered to avoid lactic acidosis such as poorly controlled diabetes, ketosis, prolonged fasting, excessive alcohol intake, hepatic insufficiency and any condition associated with hypoxia (such as decompensated cardiac failure, acute myocardial infarction).

The risk of lactic acidosis must be considered in the event of non-specific signs such as muscle cramps, digestive disorders as abdominal pain and severe asthenia. Patients should be instructed to notify these signs immediately to their physicians if they occur, notably if patients had a good tolerance to metformin before. Metformin should be discontinued, at least temporarily, until the situation is clarified. Reintroduction of metformin should then be discussed taking into account the benefit/risk ratio in an individual basis as well as renal function.

#### Diagnosis:

Lactic acidosis is characterized by acidotic dyspnea, abdominal pain and hypothermia followed by coma. Diagnostic laboratory findings are decreased blood pH, plasma lactate levels above 5 mmol/L, and an increased anion gap and lactate/pyruvate ratio. In case of lactic acidosis, the patient should be hospitalised immediately.

Physicians should alert the patients on the risk and on the symptoms of lactic acidosis.

### **Renal function:**

As metformin is excreted by the kidney, creatinine clearance (this can be estimated from serum creatinine levels by using the Cockcroft-Gault formula) or eGFR should be determined before initiating treatment and regularly thereafter:

- at least annually in patients with normal renal function,
- at least two to four times a year in patients with creatinine clearance levels at the lower limit of normal and in elderly subjects.

In case creatinine clearance CrCl is  $<45$  ml/min (eGFR  $< 45$  ml/min/1.73 m<sup>2</sup>), metformin is contraindicated.

Decreased renal function in elderly subjects is frequent and asymptomatic. Special caution should be exercised in situations where renal function may become impaired, for example in case of dehydration, or when initiating antihypertensive therapy or diuretic therapy and when starting therapy with a non-steroidal anti-inflammatory drug (NSAID).

In these cases, it is also recommended to check renal function before initiating treatment with metformin.

### **Cardiac function**

Patients with heart failure are more at risk of hypoxia and renal insufficiency. In patients with stable chronic heart failure, metformin may be used with a regular monitoring of cardiac and renal function.

For patients with acute and unstable heart failure, metformin is contraindicated.

### **Administration of iodinated contrast media:**

The intravascular administration of iodinated contrast media in radiological studies can lead to renal failure. This may induce metformin accumulation and may increase the risk for lactic acidosis. In patients with eGFR  $> 60$  mL/min/1.73m<sup>2</sup>, metformin must be discontinued prior to, or at the time of the test and not reinstated until at least 48 hours afterwards, and only after renal function has been re-evaluated and has not deteriorated further. In patients with moderate renal impairment (eGFR between 45 and 60 ml/min/1.73 m<sup>2</sup>), metformin must be discontinued 48 hours before administration of iodinated contrast media and not be reinstated until at least 48 hours afterwards and only after renal function has been re-evaluated and has not deteriorated further.

### **Surgery:**

Metformin should be discontinued 48 hours before elective surgery with general, spinal or peridural anaesthesia. Therapy may be restarted no earlier than 48 hours following surgery or resumption of oral nutrition and only if normal renal function has been established.

### **Other precautions:**

All patients should continue their diet with a regular distribution of carbohydrate intake during the day. Overweight patients should continue their energy-restricted diet.

The usual laboratory tests for diabetes monitoring should be performed regularly.

Metformin alone never causes hypoglycaemia, although caution is advised when it is used in combination with insulin or other oral antidiabetics (e.g. sulphonylureas or meglitinides).

The tablet shells may be present in the faeces. Patients should be advised that this is normal.

### **Voglibose**

The administration of Voglibose should be limited to the patients who have been definitely diagnosed as having diabetes mellitus. It should be noted that In addition to diabetes mellitus,

there are such diseases as abnormal glucose tolerance and positive urinary sugar that represent diabetes-like symptoms (renal glucosuria, senile abnormal glucose tolerance, abnormal thyroid function, etc.).

For patients who are undergoing only the basic treatment for diabetes mellitus, namely, dietary treatment and /or exercise therapy, this drug should be given only when the two-hour postprandial blood sugar is 200 mg/dL or more.

For patients who are using oral hypoglycaemic drugs or insulin preparations, in addition to dietary treatment and/or exercise therapy, a rough standard for administration of this drug is to give it when the fasting blood sugar is about 140 mg/dL or more.

During administration of Voglibose, the disease progression be closely observed with the monitoring of blood sugar at regular intervals, and careful attention should always be paid to the question of necessity for continuous administration of this drug. If its effect on postprandial blood sugar is not satisfactory even after the administration of this drug for 2 to 3 months (e.g. the reduction in the two-hour postprandial sugar level in venous plasma to 200 mg/dL or below cannot be achieved), such consideration as the change to more possible appropriate treatment should be made.

When sufficient control of the postprandial blood sugar has been attained (the two-hour postprandial sugar level reduced to 160 mg/dL or below in venous plasma), and is judged to be satisfactorily maintained only with dietary treatment and/or exercise therapy, or with additional use of oral hypoglycaemic drugs or insulin preparations, the administration of voglibose should be discontinued and the subsequent progress of disease be observed.

In administration of Voglibose, the patients should be given the sufficient explanation on hypoglycaemic symptoms and as to how they will be coped with.

### **General**

Voglibose should be administered with care in following patients:

- Patients who are receiving other antidiabetic drugs (hypoglycaemia may occur).
- Patients with a history of laparotomy or ileus (Intestinal obstruction like symptoms are liable to develop due to an increase in intestinal gas, etc.).
- Patients with chronic intestinal disease accompanied by a disturbance indigestion and absorption (the action is drug may aggravates the pathologic condition).
- Patients with Roemheld's syndrome, severe hernia, or stenosis or ulceration of the large intestinal gas, etc. (symptom may worsen due to an increase in intestinal gas, etc).
- Patients with serious hepatic dysfunction (Because of possible changes in metabolic condition, the status of blood sugar control may worsen, followed by disturbance of consciousness).

Patients with serious renal dysfunction (Because of possible changes in metabolic conditions, the status of blood sugar control may greatly vary).

- Elderly patients

## **DRUG-INTERACTION**

### **Glimepiride**

If glimepiride is taken simultaneously with certain other medicinal products, both undesired increases and decreases in the hypoglycaemic action of glimepiride can occur. For this reason, other medicinal products should only be taken with the knowledge (or at the prescription) of the doctor.

Glimepiride is metabolized by cytochrome P450 2C9 (CYP2C9). Its metabolism is known to be influenced by concomitant administration of CYP2C9 inducers (e.g. rifampicin) or inhibitors (e.g. fluconazole).

Results from an *in vivo* interaction study reported in literature show that glimepiride AUC is increased approximately 2-fold by fluconazole, one of the most potent CYP2C9 inhibitors. Based on the experience with glimepiride and with other sulfonylureas the following interactions have to be mentioned.

Potential of the blood-glucose-lowering effect and, thus, in some instances hypoglycaemia may occur when one of the following medicinal products is taken, for example:

- phenylbutazone, azapropazone and oxyfenbutazone,
- insulin and oral antidiabetic products such as metformin,
- salicylates and p-amino-salicylic acid,
- anabolic steroids and male sex hormones,
- chloramphenicol, certain long acting sulfonamides, tetracyclines, quinolone antibiotics and clarithromycin,
- coumarin anticoagulants,
- fenfluramine,
- disopyramide,
- fibrates,
- ACE inhibitors,
- fluoxetine, MAO-inhibitors,
- allopurinol, probenecid, sulfapyrazone,
- sympatholytics,
- cyclophosphamide, trophosphamide and iphosphamides,
- miconazole, fluconazole
- pentoxifylline (high dose parenteral),
- tritoqualine,

Weakening of the blood-glucose-lowering effect and, thus raised blood glucose levels may occur when one of the following medicinal products is taken, for example:

- oestrogens and progestogens,
- saluretics, thiazide diuretics,
- thyroid stimulating agents, glucocorticoids,
- phenothiazine derivatives, chlorpromazine,
- adrenaline and sympathicomimetics,
- nicotinic acid (high doses) and nicotinic acid derivatives,

- laxatives (long term use),
- phenytoin, diazoxide,
- glucagon, barbiturates and rifampicin,
- acetazolamide.

H<sub>2</sub> antagonists, beta-blockers, clonidine and reserpine may lead to either potentiation or weakening of the blood-glucose-lowering effect.

Under the influence of sympatholytic medicinal products such as beta-blockers, clonidine, guanethidine and reserpine, the signs of adrenergic counter-regulation to hypoglycaemia may be reduced or absent.

Alcohol intake may potentiate or weaken the hypoglycaemic action of glimepiride in an unpredictable fashion.

Glimepiride may either potentiate or weaken the effects of coumarin derivatives.

Colesevelam binds to glimepiride and reduces glimepiride absorption from the gastro-intestinal tract. No interaction was observed when glimepiride was taken at least 4 hours before colesevelam. Therefore, glimepiride should be administered at least 4 hours prior to colesevelam.

## **Metformin**

### Concomitant use not recommended

#### *Alcohol*

Acute alcohol intoxication is associated with an increased risk of lactic acidosis in acute alcohol intoxication, particularly in case of:

- fasting or malnutrition,
- hepatic insufficiency.

Avoid consumption of alcohol and alcohol-containing medications.

#### *Iodinated contrast media*

Intravascular administration of iodinated contrast media may lead to renal failure, resulting in metformin accumulation and an increased risk of lactic acidosis.

In patients with eGFR > 60 mL/min/1.73 m<sup>2</sup>, metformin must be discontinued prior to, or at the time of the test and not reinstated until at least 48 hours afterwards, and only after renal function has been re-evaluated and has not deteriorated further.

In patients with moderate renal impairment (eGFR between 45 and 60 ml/min/1.73 m<sup>2</sup>), metformin must be discontinued 48 hours before administration of iodinated contrast media and not be reinstated until at least 48 hours afterwards and only after renal function has been re-evaluated and has not deteriorated further.

### Combinations requiring precautions for use

*Medicinal products with intrinsic hyperglycaemic activity (e.g. glucocorticoids (systemic and local routes) and sympathomimetics).*

More frequent blood glucose monitoring may be required, especially at the beginning of treatment. If necessary, adjust the metformin dosage during therapy with the other drug and upon its discontinuation.

*Diuretics, especially loop diuretics*

They may increase the risk of lactic acidosis due to their potential to decrease renal function

## **Voglibose**

Precautions for Co-administration (Voglibose should be administered with care when Co-administered with the following drugs).

### **Antidiabetic drugs**

Derivatives of sulfonamide and sulfonurea, biguanide derivatives, insulin preparations and improving agents for insulin resistance

### **Signs, Symptoms, Treatment, Mechanisms, etc.**

It has been reported that hypoglycemia occurred in the concomitant use of Voglibose Tablets with insulin preparations or sulfonurea derivatives. Therefore, when this drug is used in combination with any of the left-listed drugs, such careful caution as starting from a lower dose should be exercised, taking into account the possible development of hypoglycaemia.

For the concomitant use of antidiabetic drugs and the drugs which enhance or diminish the hypoglycaemic action of antidiabetic drugs

- Drugs enhancing the hypoglycaemic action of antidiabetic drugs: B-blockers, salicylic acid preparations, monoamine oxidase inhibitors, fibrate derivatives for treatment of hyperlipemia, warfarin, etc.

-Drugs diminishing the hypoglycaemic action of antidiabetic drugs epinephrine, adrenocortical hormone, thyroid hormone, etc.

## **FERTILITY, PREGNANCY AND LACTATION**

### **Glimepiride**

#### Pregnancy

##### *Risk related to the diabetes*

Abnormal blood glucose levels during pregnancy are associated with a higher incidence of congenital abnormalities and perinatal mortality. So the blood glucose level must be closely monitored during pregnancy in order to avoid the teratogenic risk. The use of insulin is required under such circumstances. Patients who consider pregnancy should inform their physician.

##### *Risk related to glimepiride*

There are no adequate data from the use of glimepiride in pregnant women. Animal studies have shown reproductive toxicity which likely was related to the pharmacologic action (hypoglycaemia) of glimepiride.

Consequently, glimepiride should not be used during the whole pregnancy.

In case of treatment by glimepiride, if the patient plans to become pregnant or if a pregnancy is discovered, the treatment should be switched as soon as possible to insulin therapy.



### Lactation

The excretion in human milk is unknown. Glimepiride is excreted in rat milk. As other sulfonylureas are excreted in human milk and because there is a risk of hypoglycaemia in nursing infants, breast-feeding is advised against during treatment with glimepiride.

### **Metformin**

#### Pregnancy

Uncontrolled diabetes during pregnancy (gestational or permanent) is associated with increased risk of congenital abnormalities and perinatal mortality.

A limited amount of data from the use of metformin in pregnant women does not indicate an increased risk of congenital abnormalities. Animal studies do not indicate harmful effects with respect to pregnancy, embryonic or fetal development, parturition or postnatal development.

When the patient plans to become pregnant and during pregnancy, it is recommended that diabetes is not treated with metformin but insulin be used to maintain blood glucose levels as close to normal as possible to reduce the risk of malformations of the foetus.

#### Breast-feeding

Metformin is excreted into human breast milk. No adverse effects were observed in breastfed newborns/infants. However, as only limited data are available, breastfeeding is not recommended during metformin treatment. A decision on whether to discontinue breast-feeding should be made, taking into account the benefit of breast-feeding and the potential risk to adverse effect on the child.

#### Fertility

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately three times the maximum recommended human daily dose based on body surface area comparisons.

### **Voglibose**

#### **Paediatrics**

The safety of Voglibose in children has not been established (no clinical experience).

#### **Geriatrics**

Since the elderly have a physiological hypo function in general, the administration of Voglibose should be initiated at a lower dose (e.g. single dose of 0.1 mg).

Furthermore, this drug should be carefully administered under close observation of the course of disease conditions, with careful attention to the blood sugar level and the onset of gastrointestinal symptoms.

#### **Pregnancy**

Voglibose should be administered to pregnant women or women having possibilities of being pregnant only if the expected therapeutic benefit is thought to outweigh any possible risk [The safety of this drug In pregnant women has not been established].

### **Lactation**

It is desirable to avoid the administration of this drug to nursing mothers. However, if the administration is indispensable, nursing should be discontinued. [Animal studies (rats) have revealed a suppressive action of this drug on body weight increase in newborns, presumably due to suppression of milk production resulting from inhibition of carbohydrate absorption in mother animals].

### **EFFECTS ON ABILITY TO DRIVE AND USE MACHINES**

No studies on the effects on the ability to drive and use machines have been performed.

The patient's ability to concentrate and react may be impaired as a result of hypoglycaemia or hyperglycaemia or, for example, as a result of visual impairment. This may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating machinery).

Patients should be advised to take precautions to avoid hypoglycaemia whilst driving. This is particularly important in those who have reduced or absent awareness of the warning symptoms of hypoglycaemia or have frequent episodes of hypoglycaemia. It should be considered whether it is advisable to drive or operate machinery in these circumstances.

### **UNDESIRABLE EFFECTS**

#### **Glimepiride**

The following adverse reactions from clinical investigations were based on experience with Glimepiride and other sulfonylureas, were listed below by system organ class and in order of decreasing incidence (very common:  $\geq 1/10$ ; common:  $\geq 1/100$  to  $< 1/10$ ; uncommon:  $\geq 1/1,000$  to  $< 1/100$ ; rare:  $\geq 1/10,000$  to  $< 1/1,000$ ; very rare:  $< 1/10,000$ ), not known (cannot be estimated from the available data).

#### Blood and lymphatic system disorders

Rare: thrombocytopenia, leukopenia, granulocytopenia, agranulocytosis, erythropenia, haemolytic anaemia and pancytopenia, which are in general reversible upon discontinuation of medication. Not known: severe thrombocytopenia with platelet count less than 10,000/ $\mu$ l and thrombocytopenic purpura

#### Immune system disorders

Very rare: leukocytoclastic vasculitis, mild hypersensitivity reactions that may develop into serious reactions with dyspnoea, fall in blood pressure and sometimes shock.

Not known: cross-allergenicity with sulfonylureas, sulfonamides or related substances is possible.

#### Metabolism and nutrition disorders

Rare: hypoglycaemia.

These hypoglycaemic reactions mostly occur immediately, may be severe and are not always easy to correct. The occurrence of such reactions depends, as with other hypoglycaemic therapies, on individual factors such as dietary habits and dose.

#### Eye disorders

Not known: visual disturbances, transient, may occur especially on initiation of treatment, due to changes in blood glucose levels.

#### Gastrointestinal disorders

Very rare: nausea, vomiting, diarrhoea, abdominal distension, abdominal discomfort and abdominal pain, which seldom lead to discontinuation of therapy.

#### Hepato-biliary disorders

Not known: hepatic enzymes increased.

Very rare: hepatic function abnormal (e.g. with cholestasis and jaundice), hepatitis and hepatic failure.

#### Skin and subcutaneous tissue disorders

Not known: hypersensitivity reactions of the skin may occur as pruritus, rash, urticaria and photosensitivity.

#### Investigations

Very rare: blood sodium decrease.

### **Metformin**

In post marketing data and in controlled clinical studies, adverse event reporting in patients treated with Metformin SR was similar in nature and severity to that reported in patients treated with metformin immediate release.

During treatment initiation, the most common adverse reactions are nausea, vomiting, diarrhoea, abdominal pain and loss of appetite, which resolve spontaneously in most cases.

The following adverse reactions may occur with Glucophage SR.

Frequencies are defined as follows: very common:  $>1/10$ ; common  $\geq 1/100$ ,  $<1/10$ ; uncommon  $\geq 1/1,000$ ,  $<1/100$ ; rare  $\geq 1/10,000$ ,  $<1/1,000$ ; very rare  $<1/10,000$ .

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

#### Metabolism and nutrition disorders

Very rare:

- Lactic acidosis
- Decrease of vitamin B12 absorption with decrease of serum levels during long-term use of metformin. Consideration of such an aetiology is recommended if a patient presents with megaloblastic anaemia.

#### Nervous system disorders

*Common:*

- Taste disturbance

### Gastrointestinal disorders

*Very common:*

- Gastrointestinal disorders such as nausea, vomiting, diarrhoea, abdominal pain and loss of appetite. These undesirable effects occur most frequently during initiation of therapy and resolve spontaneously in most cases. A slow increase of the dose may also improve gastrointestinal tolerability.

### Hepatobiliary disorders

*Very rare*

- Isolated reports of liver function tests abnormalities or hepatitis resolving upon metformin discontinuation.

### Skin and subcutaneous tissue disorders

*Very rare:*

- Skin reactions such as erythema, pruritus, urticaria

### **Voglibose**

Gastrointestinal adverse effects such as diarrhoea, loose stools, abdominal pain, constipation, anorexia, nausea, vomiting, or heartburn may occur with the use of Voglibose. Also abdominal distention, increased flatus, and intestinal obstruction like symptoms due to an increase in intestinal gas, may occur with use of Voglibose.

When Voglibose is administered to patients with serious liver cirrhosis, hyperammonia may worsen with the development of constipation followed by disturbance of consciousness.

Elevation of GOT (glutamate oxaloacetate), GPT (glutamate pyruvate transaminase), LDH (lactate dehydrogenase), aGPT (aglutamate pyruvate) or alkaline phosphatase may infrequently occur. When Voglibose is used in combination with other antidiabetic drugs, hypoglycemia may occur (0.1% to <5%).

Hypersensitivity: Rash and pruritus may rarely occur. In such a case, Voglibose tablets should be discontinued.

Psychoneurologic: Headache may rarely occur.

Hematologic: Anemia; thrombocytopenia, and leucopenia may rarely occur.

Others: Numbness, edema of face, blurred vision, hot flushes, malaise, weakness, hyperkalemia, increased serum amylase, decreased HDL cholesterol, diaphoresis or alopecia, and perspiration.

### **Overdose**

#### **Glimepiride**

##### Symptoms

After ingestion of an overdose hypoglycaemia may occur, lasting from 12 to 72 hours, and may recur after an initial recovery. Symptoms may not be present for up to 24 hours after ingestion. In general observation in hospital is recommended. Nausea, vomiting and epigastric pain may occur. The hypoglycaemia may in general be accompanied by neurological symptoms like restlessness, tremor, visual disturbances, co-ordination problems, sleepiness, coma and convulsions.

Management

Treatment primarily consists of preventing absorption by inducing vomiting and then drinking water or lemonade with activated charcoal (adsorbent) and sodium-sulphate (laxative). If large quantities have been ingested, gastric lavage is indicated, followed by activated charcoal and sodium-sulphate. In case of (severe) overdose hospitalisation in an intensive care department is indicated. Start the administration of glucose as soon as possible, if necessary by a bolus intravenous injection of 50 ml of a 50% solution, followed by an infusion of a 10% solution with strict monitoring of blood glucose. Further treatment should be symptomatic.

In particular when treating hypoglycaemia due to accidental intake of Glimpiride in infants and young children, the dose of glucose given must be carefully controlled to avoid the possibility of producing dangerous hyperglycaemia. Blood glucose should be closely monitored.

### **Metformin**

Hypoglycaemia has not been seen with metformin doses of up to 85 g, although lactic acidosis has occurred in such circumstances. High overdose or concomitant risks of metformin may lead to lactic acidosis. Lactic acidosis is a medical emergency and must be treated in hospital. The most effective method to remove lactate and metformin is haemodialysis.

### **Voglibose**

The alpha-glucosidase inhibitors, acarbose, voglibose and miglitol competitively and reversibly inhibit the alpha-glucosidase enzymes (glucoamylase, sucrase, maltase and isomaltase) in the brush border in the small intestine, which delays the hydrolysis of complex carbohydrates. They appear unlikely to produce hypoglycaemia in overdose, but abdominal discomfort and diarrhoea may occur.

## **PHARMACODYNAMIC PROPERTIES**

### **Glimpiride**

#### **Pharmacodynamic properties**

Pharmacotherapeutic group: Blood glucose lowering drugs, excl. insulins: Sulfonamides, urea derivatives. ATC Code: A10B B12.

Glimpiride is an orally active hypoglycaemic substance belonging to the sulfonylurea group. It may be used in non-insulin dependent diabetes mellitus.

Glimpiride acts mainly by stimulating insulin release from pancreatic beta cells.

As with other sulfonylureas this effect is based on an increase of responsiveness of the pancreatic beta cells to the physiological glucose stimulus. In addition, glimepiride seems to have pronounced extrapancreatic effects also postulated for other sulfonylureas.

#### Insulin release

Sulfonylureas regulate insulin secretion by closing the ATP-sensitive potassium channel in the beta cell membrane. Closing the potassium channel induces depolarisation of the beta cell and results - by opening of calcium channels - in an increased influx of calcium into the cell.

This leads to insulin release through exocytosis.

Glimepiride binds with a high exchange rate to a beta cell membrane protein which is associated with the ATP-sensitive potassium channel but which is different from the usual sulfonylurea binding site.

#### Extraprocreatic activity

The extraprocreatic effects are for example an improvement of the sensitivity of the peripheral tissue for insulin and a decrease of the insulin uptake by the liver.

The uptake of glucose from blood into peripheral muscle and fat tissues occurs via special transport proteins, located in the cells membrane. The transport of glucose in these tissues is the rate limiting step in the use of glucose. Glimepiride increases very rapidly the number of active glucose transport molecules in the plasma membranes of muscle and fat cells, resulting in stimulated glucose uptake.

Glimepiride increases the activity of the glycosyl-phosphatidylinositol-specific phospholipase C which may be correlated with the drug-induced lipogenesis and glycogenesis in isolated fat and muscle cells.

Glimepiride inhibits the glucose production in the liver by increasing the intracellular concentration of fructose-2, 6-bisphosphate, which in its turn inhibits the gluconeogenesis.

#### General

In healthy persons, the minimum effective oral dose is approximately 0.6 mg. The effect of glimepiride is dose-dependent and reproducible. The physiological response to acute physical exercise, reduction of insulin secretion, is still present under glimepiride.

There was no significant difference in effect regardless of whether the medicinal product was given 30 minutes or immediately before a meal. In diabetic patients, good metabolic control over 24 hours can be achieved with a single daily dose.

Although the hydroxy metabolite of glimepiride caused a small but significant decrease in serum glucose in healthy persons, it accounts for only a minor part of the total drug effect.

#### Combination therapy with metformin

Improved metabolic control for concomitant glimepiride therapy compared to metformin alone in patients not adequately controlled with the maximum dosage of metformin has been shown in one study.

#### Combination therapy with insulin

Data for combination therapy with insulin are limited. In patients not adequately controlled with the maximum dosage of glimepiride, concomitant insulin therapy can be initiated. In two studies, the combination achieved the same improvement in metabolic control as insulin alone; however, a lower average dose of insulin was required in combination therapy.

#### Special populations

Paediatric population

An active controlled clinical trial (glimepiride up to 8 mg daily or metformin up to 2,000 mg daily) of 24 weeks duration was performed in 285 children (8-17 years of age) with type 2 diabetes.

Both glimepiride and metformin exhibited a significant decrease from baseline in HbA<sub>1c</sub> (glimepiride -0.95 (se 0.41); metformin -1.39 (se 0.40)). However, glimepiride did not achieve the criteria of non-inferiority to metformin in mean change from baseline of HbA<sub>1c</sub>. The difference between treatments was 0.44% in favour of metformin. The upper limit (1.05) of the 95% confidence interval for the difference was not below the 0.3% non-inferiority margin.

Following glimepiride treatment, there were no new safety concerns noted in children compared to adult patients with type 2 diabetes mellitus. No long-term efficacy and safety data are available in paediatric patients.

### **Pharmacokinetic properties**

#### Absorption

The bioavailability of glimepiride after oral administration is complete. Food intake has no relevant influence on absorption, only absorption rate is slightly diminished. Maximum serum concentrations ( $C_{max}$ ) are reached approx. 2.5 hours after oral intake (mean 0.3 µg/ml during multiple dosing of 4 mg daily) and there is a linear relationship between dose and both  $C_{max}$  and AUC (area under the time/concentration curve).

#### Distribution

Glimepiride has a very low distribution volume (approx. 8.8 litres) which is roughly equal to the albumin distribution space, high protein binding (>99%), and a low clearance (approx. 48 ml/min).

In animals, glimepiride is excreted in milk. Glimepiride is transferred to the placenta. Passage of the blood brain barrier is low.

#### Biotransformation and elimination

Mean dominant serum half-life, which is of relevance for the serum concentrations under multiple-dose conditions, is about 5 to 8 hours. After high doses, slightly longer half-lives were noted.

After a single dose of radiolabelled glimepiride, 58% of the radioactivity was recovered in the urine, and 35% in the faeces. No unchanged substance was detected in the urine. Two metabolites - most probably resulting from hepatic metabolism (major enzyme is CYP2C9) - were identified both in urine and faeces: the hydroxy derivative and the carboxy derivative. After oral administration of glimepiride, the terminal half-lives of these metabolites were 3 to 6 and 5 to 6 hours respectively.

Comparison of single and multiple once-daily dosing revealed no significant differences in pharmacokinetics, and the intraindividual variability was very low. There was no relevant accumulation.

#### Special populations

Pharmacokinetics were similar in males and females, as well as in young and elderly (above 65 years) patients. In patients with low creatinine clearance, there was a tendency for glimepiride clearance to increase and for average serum concentrations to decrease, most probably resulting from a more rapid elimination because of lower protein binding. Renal elimination of the two metabolites was impaired. Overall no additional risk of accumulation is to be assumed in such patients.

Pharmacokinetics in five non-diabetic patients after bile duct surgery were similar to those in healthy persons.

#### *Paediatric population*

A fed study investigating the pharmacokinetics, safety, and tolerability of a 1 mg single dose of glimepiride in 30 paediatric patients (4 children aged 10-12 years and 26 children aged 12-17 years) with type 2 diabetes showed mean  $AUC_{(0-last)}$ ,  $C_{max}$  and  $t_{1/2}$  similar to that previously observed in adults.

### **Metformin**

#### ORAL ANTI-DIABETICS

(A10BA02: Gastrointestinal tract and metabolism)

Metformin is a biguanide with antihyperglycaemic effects, lowering both basal and postprandial plasma glucose. It does not stimulate insulin secretion and therefore does not produce hypoglycaemia.

#### Mechanism of action

Metformin may act via 3 mechanisms:

- reduction of hepatic glucose production by inhibiting gluconeogenesis and glycogenolysis
- in muscle, by increasing insulin sensitivity, improving peripheral glucose uptake and utilisation
- and delay of intestinal glucose absorption.

Metformin stimulates intracellular glycogen synthesis by acting on glycogen synthase.

Metformin increases the transport capacity of all types of membrane glucose transporters (GLUT).

#### Pharmacodynamic effects

In clinical studies, the major non glycaemic effect of metformin is either weight stability or modest weight loss.

In humans, independently of its action on glycaemia, immediate release metformin has favourable effects on lipid metabolism. This has been shown at therapeutic doses in controlled, medium-term or long-term clinical studies: immediate release metformin reduces total cholesterol, LDL cholesterol and triglyceride levels. A similar action has not been demonstrated with the prolonged release formulation, possibly due to the evening administration, and an increase in triglycerides may occur.

#### Clinical efficacy:

The prospective randomised (UKPDS) study has established the long-term benefit of intensive blood glucose control in overweight type 2 diabetic patients treated with immediate release metformin as first-line therapy after diet failure. Analysis of the results for overweight patients treated with metformin after failure of diet alone showed:



- a significant reduction of the absolute risk of any diabetes-related complication in the metformin group (29.8 events/ 1000 patient-years) versus diet alone (43.3 events/ 1000 patient-years),  $p=0.0023$ , and versus the combined sulphonylurea and insulin monotherapy groups (40.1 events/ 1000 patient-years),  $p=0.0034$ .
  - a significant reduction of the absolute risk of diabetes-related mortality: metformin 7.5 events/1000 patient-years, diet alone 12.7 events/ 1000 patient-years,  $p=0.017$ ;
  - a significant reduction of the absolute risk of overall mortality: metformin 13.5 events/ 1000 patient-years versus diet alone 20.6 events/ 1000 patient-years ( $p=0.011$ ), and versus the combined sulphonylurea and insulin monotherapy groups 18.9 events/ 1000 patient-years ( $p=0.021$ );
  - a significant reduction in the absolute risk of myocardial infarction: metformin 11 events/ 1000 patient-years, diet alone 18 events/ 1000 patient-years ( $p=0.01$ )
- For metformin used as second-line therapy, in combination with a sulphonylurea, benefit regarding clinical outcome has not been shown.
- In type 1 diabetes, the combination of metformin and insulin has been used in selected patients, but the clinical benefit of this combination has not been formally established.

## **Pharmacokinetic properties**

### Absorption

After an oral dose of the prolonged release tablet, metformin absorption is significantly delayed compared to the immediate release tablet with a  $T_{max}$  at 7 hours ( $T_{max}$  for the immediate release tablet is 2.5 hours).

At steady state, similar to the immediate release formulation,  $C_{max}$  and AUC are not proportionally increased to the administered dose. The AUC after a single oral administration of 2000mg of metformin prolonged release tablets is similar to that observed after administration of 1000mg of metformin immediate release tablets b.i.d.

Intrasubject variability of  $C_{max}$  and AUC of metformin prolonged release is comparable to that observed with metformin immediate release tablets.

When the prolonged release tablet is administered in fasting conditions the AUC is decreased by 30% (both  $C_{max}$  and  $T_{max}$  are unaffected).

Mean metformin absorption from the prolonged release formulation is almost not altered by meal composition.

No accumulation is observed after repeated administration of up to 2000mg of metformin as prolonged release tablets.

Following a single oral administration of 1500 mg of Metformin SR750 mg, a mean peak plasma concentration of 1193 ng/ml is achieved with a median value of 5 hours and a range of 4 to 12 hours.

Metformin SR750 mg was shown to be bioequivalent to Metformin SR500 mg at a 1500 mg dose with respect to  $C_{max}$  and AUC in healthy fed and fasted subjects.

Following a single oral administration in the fed state of one tablet of Metformin SR1000 mg, a mean peak plasma concentration of 1214 ng/ml is achieved with a median time of 5 hours (range of 4 to 10 hours).

Metformin SR1000 mg was shown to be bioequivalent to Metformin SR500 mg at a 1000 mg dose with respect to C<sub>max</sub> and AUC in healthy fed and fasted subjects.

When the 1000 mg prolonged release tablet is administered in fed conditions the AUC is increased by 77% (C<sub>max</sub> is increased by 26% and T<sub>max</sub> is slightly prolonged by about 1 hour).

### **Distribution**

Plasma protein binding is negligible. Metformin partitions into erythrocytes. The blood peak is lower than the plasma peak and appears at approximately the same time. The red blood cells most likely represent a secondary compartment of distribution. The mean V<sub>d</sub> ranged between 63-276 L.

### **Metabolism**

Metformin is excreted unchanged in the urine. No metabolites have been identified in humans.

### **Elimination**

Renal clearance of metformin is > 400 ml/min, indicating that metformin is eliminated by glomerular filtration and tubular secretion. Following an oral dose, the apparent terminal elimination half-life is approximately 6.5 hours.

When renal function is impaired, renal clearance is decreased in proportion to that of creatinine and thus the elimination half-life is prolonged, leading to increased levels of metformin in plasma.

### Characteristics in specific groups of patients

#### Renal impairment

The available data in subjects with moderate renal insufficiency are scarce and no reliable estimation of the systemic exposure to metformin in this subgroup as compared to subjects with normal renal function could be made. Therefore, the dose adaptation should be made upon clinical efficacy/tolerability considerations.

### **Voglibose**

#### **Mechanism of Action**

Voglibose is an alpha glucosidase inhibitor which reduces intestinal absorption of starch, dextrin, and disaccharides by inhibiting the action of  $\alpha$ -glucosidase in the intestinal brush border. Inhibition of this enzyme catalyzes the decomposition of disaccharides into monosaccharides and slows the digestion and absorption of carbohydrates; the post-prandial rise in plasma glucose is blunted in both normal and diabetic subjects resulting in improvement of post prandial hyperglycemia and various disorders caused by hyperglycemia.  $\alpha$ -Glucosidase inhibitors do not stimulate insulin release and therefore do not result in hypoglycemia. These agents may be considered as monotherapy in elderly patients or in patients with predominantly post prandial hyperglycemia.  $\alpha$ -Glucosidase inhibitors are typically used in combination with other oral

antidiabetic agents and/or insulin. Voglibose should be administered at the start of a meal as it is poorly absorbed.

### **Pharmacokinetics**

Voglibose is poorly absorbed after oral doses. Plasma concentrations after oral doses have usually been undetectable. After an 80 mg dose (substantially higher than recommended dose), peak plasma levels of about 20 ng/mL were observed in 1 to 1.5 hours. When Voglibose tablets were repeatedly administered to healthy male adults (6 subjects) in a single dose of 0.2 mg, 3 times a day, for 7 consecutive days, Voglibose was not detected in plasma or urine. Similarly, when Voglibose was administered to healthy male adults (10 subjects) as a single dose of 2 mg, Voglibose was not detected in plasma or urine.

### **Distribution**

After ingestion of Voglibose (and other glucosidase inhibitors), the majority of active unchanged drug remains in the lumen of the gastrointestinal tract to exert its pharmacological activity.

### **Metabolism**

Voglibose is metabolized by intestinal enzymes and by the microbial flora.

### **Excretion**

Voglibose is excreted in the urine and feces. In a study in which a single dose of 1 mg/kg of C14-Voglibose was administered to rats, the transfer of Voglibose to the fetus and to mother's milk was observed, and the rates of excretion into urine and feces were about 5% and 98%, respectively.

### **Preclinical safety data**

#### **Glimepiride**

Preclinical effects observed occurred at exposures sufficiently in excess of the maximum human exposure as to indicate little relevance to clinical use, or were due to the pharmacodynamic action (hypoglycaemia) of the compound. This finding is based on conventional safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenicity, and reproduction toxicity studies. In the latter (covering embryotoxicity, teratogenicity and developmental toxicity), adverse effects observed were considered to be secondary to the hypoglycaemic effects induced by the compound in dams and in offspring.

#### **Metformin**

Preclinical data reveal no special hazard for humans based on conventional studies on safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity reproduction.

#### **Voglibose**

No animal studies of safety pharmacology, repeated-dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction and development have been conducted with voglibose.

### **EXPIRY DATE**

Do not use later than the date of expiry.

**STORAGE**

Store in a cool, dry place  
Protect from light.

**PRESENTATION**

Trivoglitor Forte 1 and Trivoglitor Forte 2 are available in blister pack of 10 tablets.

**MARKETED BY**

TORRENT PHARMACEUTICALS LTD.

Torrent House, Off Ashram Road,  
Ahmedabad-380 009, INDIA

**IN/TRIVOGLITOR FORTE 0.3,1,2,500mg/JAN-2016/01/PI**