

SERIVA

For the use only of a Dermatologist or a Hospital or a Laboratory

Abbreviated Prescribing information for Seriva lotion (sertaconazole nitrate lotion, 2% w/v) [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Sertaconazole nitrate is similar to that of other imidazoles, in that it inhibits fungal biosynthesis of ergosterol, resulting in disorganization of the fungal plasma membrane and altered permeability.

INDICATION: It is indicated for the topical treatment of superficial fungal infections of skin including tenia pedis.

DOSAGE AND ADMINISTRATION: In the treatment of superficial fungal infections of skin, it should be applied twice daily for 2 weeks. In the treatment of interdigital tenia pedis, it should be applied twice daily for 4 weeks.

DIRECTION FOR USE: Sufficient sertaconazole nitrate lotion, 2%, should be applied to cover both the affected areas and the immediately surrounding healthy skin. For topical use only, not for oral, ophthalmic or intravaginal use.

CONTRAINDICATION: Contraindicated in patients who have a known or suspected sensitivity to sertaconazole nitrate or any of its components or to other imidazoles.

WARNINGS & PRECAUTIONS: It is not indicated for ophthalmic, oral or intravaginal use. If irritation or sensitivity develops with the use, treatment should be discontinued and appropriate therapy instituted. Information for Patients: The patient should be instructed to: The hands should be washed after applying the medication, avoid contact with the eyes, nose, mouth and other mucous membranes it is for external use only, dry the affected area thoroughly before application, use the medication for the full treatment time recommended by the physician, even though symptoms may have improved, notify the physician if there is no improvement after the end of the prescribed treatment period, or sooner, if the condition worsens and inform the physician if the area of application shows signs of increased irritation, redness, itching, burning, blistering, swelling or oozing and avoid the use of occlusive dressings unless otherwise directed by the physician.

DRUG INTERACTION: Potential interactions between sertaconazole and other drugs or laboratory tests have not been systematically evaluated.

ADVERSE REACTIONS: Contact dermatitis, dry skin, burning skin, application site reaction and skin tenderness, erythematous reaction, pruritus, vesiculation, desquamation and hyperpigmentation.

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.
Torrent House, Off Ashram Road,
Ahmedabad-380 009, INDIA

IN/SERIVA LOTION 2%/Mar-18/01/AbPI
(Additional information is available on request)