

UNIENZYME

For the use of a Registered Medical Practitioner or Hospital or a Laboratory only. Abbreviated Prescribing information for Unienzyme (Fungal Diastase, Papain & Activated Charcoal Tablets) [Please refer the complete prescribing information available at www.torrentpharma.com].

PHARMACOLOGICAL PROPERTIES: a Unienzyme is a combination of Papain, Fungal Diastase and Activated charcoal. Papain is an enzyme obtained from the juice of the unripe fruit of *Carica papaya* Linn that digests protein and catalyses the breakdown of proteins by hydrolysis. It is sparingly soluble in water. Fungal Diastase is a potent, dependable and powerful enzyme complex obtained from *Aspergillus oryzae*. Fungal Diastase is a starch-degrading enzyme. It helps in digestion of starch and carbohydrates from the diet into simple polysaccharides and sugar.

Unienzyme tablets are black coloured, oval shaped, sugar coated tablets, having “Unienzyme” printed in white along one side. The excipients used are Colloidal Silicon Dioxide, Microcrystalline Cellulose, Dibasic Calcium Phosphate Dihydrate, Lactose Monohydrate, Gum Acacia, Gelatin, Sodium Benzoate, Talc, Magnesium Stearate, Shellac, Isopropyl Alcohol, Castor Oil, Calcium Carbonate, Sugar, Vegetable Charcoal Fine, Carnauba Wax, and Edible Ink White.

DOSAGE AND ADMINISTRATION:

As directed by the physician.

Method of Administration- Oral use.

THERAPEUTIC INDICATION

This is indicated as symptomatic relief of functional dyspepsia and other gastrointestinal disorders which usually manifest with symptoms of bloating, belching, flatulence, abdominal discomfort or feeling of abdominal fullness

CONTRAINDICATION:

Unienzyme is contraindicated in patients with known hypersensitivity to any of the ingredients of the formulation.

SPECIAL WARNINGS AND PRECAUTIONS FOR USE

History of the allergy, impairment of hepatic or renal function, G6-PD deficiency. To get periodic blood check & liver function tests to be done.

Unienzyme is used with caution in patients with an allergy to fig and kiwi. Do not exceed the stated dose. Check the expiry date on the pack and do not use after expiry.

DRUGS INTERACTIONS

Not available

ADVERSE REACTIONS: Prolonged use in patients may cause nausea, Skin Problems

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Indrad-382 721, Dist. Mehsana, INDIA.

IN/UNIENZYME 100, 60, 75 mg /May-2019/01/ABPI

(Additional information is available on request)