

U.V.A. CEF DRY SYRUP

For the use of a Registered Medical Practitioner or Hospital or a Laboratory only.
Abbreviated Prescribing information for U.V.A. CEF DRY SYRUP (Cefpodoxime Proxetil
Oral Suspension)

[Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Cefpodoxime is a bactericidal agent that acts by inhibition of bacterial cell wall synthesis. Cefpodoxime has activity in the presence of some beta-lactamases, both penicillinases and cephalosporinases, of Gram-negative and Gram-positive bacteria.

DOSAGE AND ADMINISTRATION:

This product is for Paediatric use only

Dosage should be taken as directed by your Physician.

Direction for reconstitution: To make 30ml of suspension: Shake the bottle well to loosen the powder. Use enclosed sterile water supplied with this pack to reconstitute the powder up to the mark on the bottle. Discard remaining sterile water. After reconstitution keep the bottle in a refrigerator when not in use and use the reconstituted suspension within 14 days.

CONTRAINDICATION: Known hypersensitivity to cephalosporin antibiotics or to any of the excipients.

WARNINGS & PRECAUTIONS: UVA CEF Dry Syrup, in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug resistant bacteria.

DRUG INTERACTION: It can interact with Antacids, Probenecid, Nephrotoxic drugs, Certain Drug/Laboratory Test Interactions.

ADVERSE REACTIONS: Diarrhoea, Diaper rash/Fungal skin rash, Other skin rashes, Vomiting, Localized abdominal pain, abdominal cramp, headache, monilia, generalized abdominal pain, asthenia, fever, fungal infection, Nausea, monilia, anorexia, dry mouth, stomatitis, pseudomembranous colitis, Thrombocytopenia, positive direct Coombs' test, eosinophilia, leukocytosis, leukopenia, prolonged partial thromboplastin time, thrombocytopenic purpura, Increased SGPT, Myalgia, Hallucination, hyperkinesia, nervousness, somnolence, Epistaxis, rhinitis, Skin moniliasis, urticaria, fungal dermatitis, acne, exfoliative dermatitis, maculopapular rash and Taste perversion.

MARKETED BY:

TORRENT PHARMACEUTICALS LTD.

Torrent House, Off Ashram Road,

Ahmedabad-380 009, INDIA

IN/ U.V.A. CEF DRY SYRUP 50mg, 100mg/Aug-2020/01/AbPI

(Additional information is available on request)