



Dispense with Patient Information available at:
<https://torrentpharma.com/pi/usa/products/>

PATIENT INFORMATION
Sildenafil (sil DEN a fil) Tablets USP

What is the most important information I should know about Sildenafil tablets?
Never take Sildenafil tablets with any nitrate or guanylate cyclase stimulator medicines.

- Your blood pressure could drop quickly to an unsafe level. Nitrates include:
 - Medicines that treat chest pain (angina)
 - Nitroglycerin in any form including tablets, patches, sprays, and ointments
 - Isosorbide mononitrate or dinitrate
 - Street drugs called “poppers” (amyl nitrate, butyl nitrate or nitrite)

Guanylate cyclase stimulators include:

- Riociguat, a medicine that treats pulmonary arterial hypertension and chronic thromboembolic pulmonary hypertension.

Ask your healthcare provider or pharmacist if you are not sure if you are taking a nitrate or a guanylate cyclase stimulator medicine.

See “**What are the possible side effects of Sildenafil tablets?**” for more information about side effects.

What are Sildenafil tablets?

Sildenafil tablets are prescription medicine used to treat pulmonary arterial hypertension (PAH). PAH is a type of high blood pressure in the arteries of your lungs. Sildenafil tablets may be used in:

- adults to improve your ability to exercise and help slow down the worsening of your physical condition.

It is not known if Sildenafil tablets are safe and effective in children younger than 1 year of age.

Do not take Sildenafil tablets if you:

- take medicines called nitrates.
- take riociguat, a guanylate cyclase stimulator medicine.
- are allergic to sildenafil or any of the ingredients in Sildenafil tablets. See the end of this leaflet for a complete list of ingredients in Sildenafil tablets.

Before taking Sildenafil tablets, tell your healthcare provider about all of your medical conditions, including if you:

- have low blood pressure
- have heart problems
- have pulmonary veno-occlusive disease (PVOD)

- have bleeding problems or a stomach (peptic) ulcer. It is not known if Sildenafil tablets are safe in people with bleeding problems or who have a stomach ulcer.
- have an eye problem called retinitis pigmentosa
- have ever had sudden loss of vision in one or both eyes, including an eye problem called non-arteritic anterior ischemic optic neuropathy (NAION)
- have ever had hearing problems such as ringing in the ears, dizziness, or loss of hearing
- have a deformed penis shape or Peyronie’s disease
- have any blood cell problems such as sickle cell anemia
- are pregnant or plan to become pregnant. It is not known if sildenafil tablets will harm your unborn baby.
- are breastfeeding or plan to breastfeed. Sildenafil citrate passes into your breast milk. It is not known if it can harm your baby. Talk with your healthcare provider about the best way to feed your baby during treatment with sildenafil tablets.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Sildenafil tablets and certain other medicines may affect each other and can cause side effects.

Especially tell your healthcare provider if you take:

nitrates or guanylate cyclase stimulators. See “**What is the most important information I should know about sildenafil tablets?**”

- medicines to treat high blood pressure
- medicines for erectile dysfunction (impotence). sildenafil tablets contain sildenafil, which is the same medicine found in another medicine called VIAGRA®. VIAGRA is used for the treatment of erectile dysfunction. **Do not take VIAGRA or other PDE-5 inhibitors during treatment with sildenafil tablets.**

Ask your healthcare provider or pharmacist for a list of these medicines if you are not sure.

Know the medicines you take. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.

How should I take Sildenafil tablets?

- Take or give Sildenafil tablets exactly as your healthcare provider tells you.
- Your healthcare provider may change your dose of Sildenafil tablets as needed. Do not

change your dose or stop taking Sildenafil tablets without talking to your healthcare provider.

- Take your prescribed dose of Sildenafil tablets 3 times a day.
- If you take too much Sildenafil tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

What are the possible side effects of Sildenafil tablets?

Sildenafil tablets may cause serious side effects, including:

- See “**What is the most important information I should know about Sildenafil tablets?**”
- Decreased blood pressure.** Sildenafil tablets may cause low blood pressure that last for a short time. If you take medicines to treat high blood pressure, your healthcare provider should monitor your blood pressure during treatment with Sildenafil tablets.
- Decreased eyesight or permanent loss of vision in one or both eyes** can be a sign of non-arteritic anterior ischemic optic neuropathy (NAION). Most people who develop NAION have certain risk factors. You can ask your healthcare provider if you have questions about risk factors for NAION. If you notice a sudden decrease or loss of vision in one or both eyes during treatment with Sildenafil tablets, contact your healthcare provider right away.
- Sudden decrease or loss of hearing**, sometimes with ringing in the ears and dizziness. If you notice a sudden decrease or loss of hearing during treatment with Sildenafil tablets, contact your healthcare provider right away.
- In men, an erection that lasts for more than 4 hours (priapism).** If you have an erection, with or without pain, that lasts more than 4 hours, contact your healthcare provider or get emergency medical help right away. A painful erection that lasts more than 6 hours must be treated right away or you can have lasting damage to your penis, including the inability to have erections.

The most common side effects of Sildenafil tablets in adults include:

- nosebleeds
- muscle aches and pain
- headache
- back pain
- upset stomach
- diarrhea
- getting red or hot in the face (flushing)
- arm or leg pain

These are not all the possible side effects of Sildenafil tablets.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Sildenafil tablets?

- Store Sildenafil tablets at room temperature between 68°F to 77°F (20°C to 25°C).

Keep Sildenafil tablets and all medicines out of the reach of children.

General information about the safe and effective use of Sildenafil tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use Sildenafil tablets for a condition for which it was not prescribed. Do not give Sildenafil tablets to other people, even if they have the same symptoms that you have. It may harm them. You can ask your healthcare provider or pharmacist for information about Sildenafil tablets that is written for health professionals.

What are the ingredients in Sildenafil tablets?

Active ingredients: sildenafil citrate, USP

Inactive ingredients:

Sildenafil tablets: croscarmellose sodium, dibasic calcium phosphate anhydrous, hypromellose, microcrystalline cellulose, sodium stearyl fumarate, titanium dioxide and triacetin.

Pediatric use information is approved for Viatris Specialty LLC’s, REVATIO (sildenafil) tablets. However, due to Viatris Specialty LLC’s marketing exclusivity rights, this drug product is not labeled with that information.

Trademarks are the property of their respective owners.

For more information call Torrent Pharma Inc. at 1-800-912-9561.



Manufactured by:
Torrent Pharmaceuticals LTD., India.

Manufactured for:
Torrent Pharma INC.
Basking Ridge, NJ 07920.

8098158 Revised: November 2024

This Patient Information has been approved by the U.S. Food and Drug Administration.



PRODUCT NAME :	SILDENAFIL TAB 20MG USP	COUNTRY : US	LOCATION : Indrad/Dahej	Supersedes A/W No.:
ITEM / PACK :	Medication Guide	NO. OF COLORS: 1	REMARK :	V. No. : 01
DESIGN STYLE :	Front / Back Side	PANTONE SHADE NOS.:	SUBSTRATE :	
CODE :	8098158	Black	Activities	Department
DIMENSIONS (MM) :	150 x 250		Name	Signature
ART WORK SIZE :	S/S		Prepared By	Pkg. Dev.
DATE :	20-11-2024	Font Size 10 pt	Reviewed By	Pkg. Dev.
			Approved By	Quality



5076