

PREGALIN M SR

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

Abbreviated Prescribing information for Pregalin M SR (Pregabalin 75mg,150mg Sustained Release And Methylcobalamin 1500mcg Tablets) [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: *Pregabalin*: It is a structural derivative of the inhibitory neurotransmitter gammaaminobutyric acid (GABA). *Methylcobalamin* is required for synthesis of thymidine from deoxyuridine and promotes the synthesis of DNA and RNA.

INDICATION: It is indicated for painful neuropathies such as diabetic neuropathy, postherpetic neuralgia, trigeminal neuralgia, following to nerve entrapment as in prolapsed intervertebral disc, complex regional pain syndrome, infective neuropathies, HIV sensory neuropathy, alcoholic neuropathy, nutritional neuropathies and phantom limb pain.

DOSAGE AND ADMINISTRATION: It should be taken once daily with/without food. Maximum dose can be 300 mg/day and dosage above 300 mg/day must be reserved only for those having ongoing pain, and tolerating 300mg pregabalin per day. While discontinuing, it must be gradually reduced over 1 week. Do not crush or chew the tablet. Take whole tablet with water.

CONTRAINDICATION: In patients who are hypersensitive to the drug or to any of its ingredients.

WARNINGS & PRECAUTIONS: It should be given cautiously in heart failure, diabetic patients, Leber's disease and renal dysfunction. If the potential benefit justifies the probable risks, it should be taken during pregnancy and lactation. It can manifest of central nervous system depression and cause vision disturbance. It is not to be taken along with alcohol, or be combined with other central nervous depressants.

DRUG INTERACTIONS: Gabapentin may slightly decrease the rate of absorption of pregabalin.

ADVERSE REACTIONS: Weight gain, edema, skin ulceration, dizziness, somnolence, dry mouth, edema, blurred vision, weight gain, and "thinking abnormal" (primarily difficulty with concentration/attention). Asthenia, ataxia, abnormal gait, paresthesias, diplopia, blurred vision, visual field defect, nystagmus, nausea, tremor, vertigo, headache, nervousness, confusion and edema. Flu syndrome, dry mouth, myasthenia, bronchitis, diplopia, urinary incontinence, allergic reaction, suicide attempt, cellulitis, pelvic pain, deep thrombophlebitis, heart failure, gastroenteritis, gastrointestinal hemorrhage, increased appetite, glucose tolerance decreased, ecchymosis, anxiety, libido decreased, apnea, stevens-johnson syndrome, lung fibrosis, anorgasmia, impotence, nausea, vomiting and diarrhea. Pulmonary edema and congestive heart failure, mild transient diarrhea, itching; transitory exanthema and feeling of swelling of entire body.

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Torrent House, Off Ashram Road,
Ahmedabad-380 009, INDIA

IN/ PREGALIN M SR/75mg,150mg,1500mcg/May-15/01/AbPI

(Additional information is available on request)