

STROLIN P

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only

Abbreviated Prescribing information for STROLIN P (STROLIN P 400 (Citicoline 500 mg, Piracetam 400 mg) STROLIN P 800 (Citicoline 500 mg, Piracetam 800 mg)) [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: **Citicoline:** A naturally occurring endogenous nucleoside, is an intermediate compound in the major pathway for the biosynthesis of the structural phospholipids of cell membranes, including neurons. **Piracetam:** Piracetam is a nootropic, which is claimed to enhance cognition and memory, slow down brain aging, increase blood flow and oxygen to the brain, aid stroke recovery, and improve Alzheimer's, Down syndrome, dementia, and dyslexia, among others. **INDICATION:** Fixed-dose combination of Citicoline and Piracetam is indicated in the management of acute stroke.

DOSAGE AND ADMINISTRATION: One tablet 2 times daily

CONTRAINDICATION: Contraindicated in patients with severe renal impairment, hepatic impairment and to those under 16 years of age. It is also contraindicated in patients with cerebral haemorrhage, suffering from Huntington's Chorea and in those with hypersensitivity to any of the ingredients, other pyrrolidone derivatives or any of the excipients.

WARNINGS & PRECAUTIONS: Due to the effect of piracetam on platelet aggregation, caution is recommended in patients with severe haemorrhage, patients at risk of bleeding such as gastrointestinal ulcer, patients with underlying disorders of haemostasis, patients with history of haemorrhagic CVA, patients undergoing major surgery including dental surgery, and patients using anticoagulants or platelet antiaggregant drugs. Piracetam is eliminated via the kidneys and care should thus be taken in cases of renal insufficiency. For long-term treatment in the elderly, regular evaluation of the creatinine clearance is required to allow dosage adaptation if needed. Abrupt discontinuation of treatment should be avoided as this may induce myoclonic or generalised seizures in some myoclonic patients

DRUG INTERACTIONS: **Thyroid hormones:** Confusion, irritability and sleep disorder have been reported during concomitant treatment with thyroid extract (T3 + T4). **Acenocoumarol:** compared with the effects of acenocoumarol alone, the addition of piracetam significantly decreased platelet aggregation, β -thromboglobulin release, levels of fibrinogen and von Willebrand's factors and whole blood and plasma viscosity.

ADVERSE REACTIONS: Numbness, headache, tingling sensations, abdominal discomfort, diarrhea, haemorrhagic disorder, anaphylactoid reaction, hypersensitivity, nervousness, agitation, anxiety, confusion, hallucination, hyperkinesias, ataxia, balance impaired, epilepsy aggravated, insomnia, vertigo, abdominal pain, nausea, vomiting, angioneurotic oedema, dermatitis, pruritus, urticaria, weight increased.

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Torrent House, Off Ashram Road,

Ahmedabad-380 009, INDIA

IN/ STROLIN P/MAR 2015/01/AbPI

(Additional information is available on request)

