

PREGEB

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory Only

Abbreviated Prescribing information for **Pregeb** [Pregabalin 75mg and 150mg Capsules] [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Pregabalin binds with high affinity to the alpha2-delta site (an auxiliary subunit of voltage gated calcium channels) in central nervous system tissues and it is a structural analogue of gamma amino butyric acid (GABA) and thus produces anti-epileptic effect and relief against neuropathic pain.

INDICATION: For the treatment of neuropathic pain in adults.

DOSAGE AND ADMINISTRATION: Pregabalin taken orally with or without food. Dose range is 150 to 600 mg per day given in either two or three divided doses. **Neuropathic pain:** starting dose of 150 mg per day given as two or three divided doses, the dose may be increased to 300 mg per day after an interval of 3 to 7 days, and if needed, to a maximum dose of 600 mg per day after an additional 7-day interval. Discontinuation is recommended gradually over a minimum of 1 week.

CONTRAINDICATION: Contraindicated in patients with known hypersensitivity to pregabalin or any components of this product.

WARNINGS & PRECAUTIONS: Use cautiously or stop the pregabalin in patient with withdrawal of antiepileptic drugs, tumorigenic potential, angioedema, dizziness and somnolence, blurred vision, weight gain, peripheral edema, creatine kinase elevations, decreased platelet count, PR interval prolongation, suicidal ideation and behavior, constipation, encephalopathy and Abrupt or rapid discontinuation of pregabalin in some patients reported insomnia, nausea, headache, and diarrhea.

DRUG INTERACTIONS: No specific drug interactions.

ADVERSE REACTIONS: dizziness, Somnolence, asthenia, confusion, and peripheral edema, asthenia, ataxia, and abnormal gait, infection, headache, pain, accidental injury, vertigo, dry mouth, constipation, vomiting, weight gain, myasthenia, bronchitis, blurry vision, diplopia, abnormal vision, urinary incontinence, abdominal pain, allergic reaction, fever, gastroenteritis, increased appetite, ecchymosis, arthralgia, leg cramps, myalgia, anxiety, hypertonia, hypesthesia, libido decreased, nystagmus, paresthesia, stupor, twitching, pruritus, conjunctivitis, diplopia, otitis media, tinnitus, anorgasmia, impotence, urinary frequency, muscle cramp, arthralgia, back pain, pain in limb, cervical spasm, oedema peripheral, oedema, gait abnormal, fall, feeling drunk, feeling abnormal, fatigue and disorientation

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Torrent House, Off Ashram Road,
Ahmedabad-380 009, INDIA

IN/ PREGEB 75,150mg/Aug-15/01/ABPI

(Additional information is available on request)