CARNISURE

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory Only

Abbreviated Prescribing information for CARNISURETM [Levocarnitine 500mg Tablets and Syrup] [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Levocarnitine is a naturally occuring substance required in mammalian energy metabolism. It has been shown to facilitate long-chain fatty acid entry into cellular mitochondria, therefore delivering substrate for oxidation & subsequent energy production.

INDICATION: It is indicated in the treatment of primary systemic carnitine deficiency. It is useful in the treatment of cardiomyopathy, skeletal myopathy, ischaemic cardiopathy, long-term haemodialysis, oligospermia, hereditary disorders affecting fatty acid metabolism, acute neonatal crisis, failure to thrive, in preterm infants and along with valproate therapy.

DOSAGE AND ADMINISTRATION: *Adults:* The recommended oral dosage for adults is 990 mg two or three times a day and increased slowly, while assessing tolerance & therapeutic response. With the oral solution the recommended dose is 10-30 mg/kg for a 50 kg subject. *Infants & Children:* The dose recommended is 50-100 mg/kg/day.

CONTRAINDICATION: None known.

WARNINGS & PRECAUTIONS: High dose not recommended in patients with severely renal compromised patient/ESRD patients as major metabolite trimethylamine-N-oxide (TMAO) will accumulate. Inefficient removal of TMA may result in development of "fishy odor" syndrome.

DRUG INTERACTIONS: None known.

ADVERSE REACTIONS: Various mild gastrointestinal complaints have been reported during the long-term administration of oral levocarnitine, these include transient nausea, vomiting, abdominal cramps and diarrhoea.

MARKETED BY:



TORRENT PHARMACEUTICALS LTD. Torrent House, Off Ashram Road, Ahmedabad-380 009, INDIA

IN/CARNISURE/Feb-2015/01/AbPI

(Additional information is available on request)