

## CHYMORAL FORTE / CHYMORAL FORTE DS

**For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only**

Abbreviated Prescribing information for CHYMORAL FORTE/ CHYMORAL FORTE – DS (Trypsin-Chymotrypsin Tablets) [Please refer the complete prescribing information available at [www.torrentpharma.com](http://www.torrentpharma.com)]

**PHARMACOLOGICAL PROPERTIES:** Trypsin and Chymotrypsin are proteolytic enzymes which have anti-inflammatory properties by following ways: Fibrinolytic activity, Reduction in Plasmin Inhibitor levels, Release of Intestinal Plasminogen activators, Increased Microcirculation and also Smoothens process of digestion.

**INDICATION:** Acute or chronic osteoarthritis, rheumatoid arthritis and spondyloarthritis, spondylosis and other ortho-degenerative disorders, pain management, Post-operative wounds, Oedema, Haematoma, Prevention of inflammation of the stitching, Pelvic Inflammatory Disease, Caesarean section, Episiotomy, Hysterectomy, Tooth Extraction, Peri apical abscess, Maxillofacial surgery, Ocular Trauma such as Macular edema, Black eye, Hyphema, Uveal tract inflammation, Subconjunctival hemorrhage, Auricular Septal hematoma, Nasal fractures, Parapharyngeal abscess, Post-traumatic edema, soft tissue injury, Fractures & dislocation, Sports injury, Sprains & Strains, Intervertebral disc herniation, and Extra-ocular trauma.

**DOSAGE AND ADMINISTRATION:** As directed by the Physician. The dose and dose frequency will be decided under the supervision of qualified physician.

**CONTRAINDICATION:** Adjunctive therapy in management of inflammatory edema. Hypersensitivity to Chymoral ingredients or enzymes, Chymoral is contraindicated in patients with severe liver, kidney impairment, peptic ulcer, high vitreous pressure, and hypersensitivity.

**WARNINGS & PRECAUTIONS:** When used in the eye, it can cause increase in pressure in the eye and other eye conditions such as uveitis, paralysis of the iris, and keratitis. Not be employed in patients with severe hepatic insufficiency or renal damage or irregularities of blood clotting mechanism. To be used with caution during Lactation, or in the elderly, children, pregnancy and patients with irregularities of blood clotting mechanism.

**DRUG INTERACTION:** Systemic proteases may increase the effectiveness of herbal supplements. Chymotrypsin is also known to interact with alcohol. Antibiotics: increase in the levels of the semi synthetic penicillin antibiotics and interact with chloramphenicol. Anticoagulants: CHYMORAL FORTE – DS should not be administered concurrently with anti-coagulants.

**ADVERSE REACTIONS:** Hepatic damage and necrosis may precipitate arrhythmias, shivering during recovery. allergic reaction when taken by mouth having Symptoms like itching, shortness of breath, swelling of the lips or throat, shock, loss of consciousness, and death.

### MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Torrent house, Off Ashram road,  
Ahmedabad - 380009, INDIA

IN/ CHYMORAL FORTE/CHYMORAL FORTE – DS/FEB 2015/01/AbPI  
(Additional information is available on request)