

CYCLOSPASMOL

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only

Abbreviated Prescribing information for CYCLOSPASMOL (Cyclandelate 200 mg and 400 mg capsule) [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Cyclandelate produces peripheral vasodilation by a direct effect on vascular smooth muscle. Pharmacological action may be due to calcium-channel antagonism.

INDICATION: Peripheral vascular disease: Atherosclerosis, Vasospastic states e.g. Raynaud's disease, Buerger's disease, Cerebrovascular disease, Transient Ischaemic attacks initiated by vasospasm, hypertensive encephalopathy; neurological defects such as paresia, hemiplegia, aphasia and cerebral atherosclerosis with disturbance of concentration, disorientation, vertigo, insomnia, confusion, fatigue, loss of memory and disturbance in behaviour. One must remember that these diseases can manifest themselves in several ways, depending on the localization.

DOSAGE AND ADMINISTRATION: 800 to 1600 mg (4 to 8 Capsules) daily or as directed by the Physician. It is given orally in an initial dosage of up to 2 g daily in divided doses; a usual maintenance dose is 0.8 to 1.2 g daily.

CONTRAINDICATION: Acute phase of cerebrovascular accidents.

WARNINGS & PRECAUTIONS: Patients with severe obliterative coronary artery disease or cerebrovascular disease.

ADVERSE REACTIONS: Flushes, GI upset, nausea, tingling, tachycardia, sweating, dizziness, headache.

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Torrent house, Off Ashram road,

Ahmedabad - 380009, INDIA

IN/ CYCLOSPASMOL /MAR 2015/01/AbPI

(Additional information is available on request)