

PREXAN

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only

Abbreviated Prescribing information for PREXAN [Ferrous fumarate equivalent to elemental iron 60mg, folic acid 1mg, cyanocobalamin 7.5mcg, copper gluconate equivalent to elemental copper 1mg, docusate sodium 50mg] [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Multivitamin and multimineral. PREXAN is a haematinic that contains iron salts (ferrous fumarate), folic acid, cyanocobalamin and docusate sodium.

INDICATION: Iron deficiency: For the prevention and treatment of iron deficiency and iron deficiency anemia. Iron supplement: As a dietary supplement for iron.

DOSAGE AND ADMINISTRATION: The usual prophylactic dose in adults is about 60mg of elemental iron daily. The usual adult treatment dose is 100 to 200mg of supplemental iron daily. 1-2 capsules per day depending on the nutritional status of the patients or as directed by the physician. Iron Supplementation: In pregnancy 15 to 30mg elemental iron daily (not taken with meals) should be adequate to meet the daily requirement of the last 2 trimesters.

WARNINGS & PRECAUTIONS: Caution is advised in patient with pernicious anemia and other megaloblastic anemia, chronic iron intake, renal impairment and hepatic impairment.

DRUG INTERACTION: Cimetidine, fluroquinolones, pancreatin or pancreatic lipase, tetracyclines, grain breads and cereals (contain phytic acid and dietary fiber), eggs and milk, coffee and tea.

ADVERSE REACTIONS: Nausea, vomiting, constipation, diarrhea, stain the teeth, anorexia, heartburn and allergic sensitizations.

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Torrent house, Off Ashram road,

Ahmedabad - 380009, INDIA

IN/PREXAN/MARCH 2015/01/AbPI

(Additional information is available on request)