COLONISE MV

For the use of a Registered Medical Practitioner or Hospital or a Laboratory only
Abbreviated Prescribing information for COLONISE MV [Pre and Probiotic, Vitamins and Minerals
Capsules] [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Colonise MV contains a pre & probiotic, vitamins and minerals combination which restores the gastrointestinal microflora, thus minimizes gastrointestinal infection, regulates bowel movements and supports gastrointestinal health. Streptococcus faecalis, Clostridium butyricum and Bacillus mesentericus are a probiotic. Vitamins and minerals corrects vitamin deficiency and probiotics which are defined as live microorganisms, including Lactobacillus species, Bifidobacterium species and yeasts beneficially affect the host upon ingestion by improving the balance of the intestinal microflora.

INDICATION: It is used to correct vitamins and minerals deficiency and also helps to replace the lost beneficial bacteria in adult patients. It is for Therapeutic use only.

DOSAGE AND ADMINISTRATION: One Capsule twice a day or as directed by the physician for optimum gastrointestinal health as it restores intestinal microflora and corrects vitamin and mineral deficiency.

CONTRAINDICATION: It is contraindicated in the patients who are having hypersensitivity to active constituents or any of the formulation ingredients.

WARNINGS & PRECAUTIONS: There are no known hazards to the use of colonise MV in pregnancy; supplements during pregnancy are often beneficial. But it should be used under physician's concern only.

DRUG INTERACTIONS: Absorption from the gastrointestinal tract may be reduced by neomycin, aminosalicylic acid, histamine H2-antagonists, omeprazole, and colchicine. Serum concentrations may be decreased by use of oral contraceptives, chloramphenicol and co-trimoxazole may interfere with folate metabolism, Sulfasalazine - can reduce the absorption of folic acid. Folic acid may interfere with the toxic and therapeutic effects of Methotrexate. The absorption of zinc may be reduced by iron supplements, penicillamine, phosphorus containing preparations, and tetracyclines. Zinc supplements reduce the absorption of copper, fluoroquinolones, iron, penicillamine and tetracyclines.

ADVERSE REACTIONS: Mild transient diarrhea, rarely itching, transitory exanthema, Anorexia, nausea, abdominal distention and flatulence, constipation, Allergic reactions, comprising erythema, rash, pruritus, urticaria, dyspnea and anaphylactic reactions (including shock).

MARKETED BY:



TORRENT PHARMACEUTICALS LTD. Torrent House, Off Ashram Road, Ahmedabad-380 009, INDIA

IN/ COLONISE MV /JUL-2017/01/ABPI

(Additional information is available on request)