UNIENZYME LIQUID

For the use of a Registered Medical Practitioner or Hospital or a Laboratory only. Abbreviated Prescribing information for UNIENZYME LIQUID (Digestive Enzymes)

[Please refer the complete prescribing information available at <u>www.torrentpharma.com</u>].

PHARMACOLOGICAL PROPERTIES: Fungal Diastase- An enzyme catalyzing the hydrolysis of linkages of polysaccharides such as starch. Best quality of amylase (diastase) is available from Aspergillus oryzae. The major amyloytic activity of diastase (Fungal Diastase) is on starch as alpha amylase and S-amylase (Saccharifying amylase), and has some proteolytic activity in addition, it also contains lipase, celluose, pectinase and several other enzymes making it a rich source of enzyme complex which splits carbohydrates, proteins, fats and cellulosic constituents of food in their assimilable forms and thus greatly enhances digestive capacity. Amylase has been used as a digestant.

Pepsin- Pepsin is a proteolytic enzyme which is secreted by the stomach and controls the degradation of proteins into proteoses and peptones. It hydrolysis polypeptides. Pepsin is a substance containing a proteolytic enzyme present in the gastric juice of animals and obtained from the membrane of the stomach of certain animals commonly used for food.

DOSAGE AND ADMINISTRATION: Adults: 5ml (1 teaspoonful) immediately after meal or as directed by the physician Children: Half the adult dose. Shake well before use.

Method of Administration- Oral use.

CONTRAINDICATION: Hypersensitivity to any of the ingredients.

ADVERSE REACTIONS: Prolonged use in patients ingesting low phosphate diet may cause osteomalacia, nausea, vomiting, constipation.

MARKETED BY:

TORRENT PHARMACEUTICALS LTD. Torrent House, Off Ashram Road, Ahmedabad-380 009, INDIA

IN/ UNIENZYME LIQUID/APR-2020/01/ABPI

(Additional information is available on request)