

DOMSTAL DT

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only

Abbreviated Prescribing information for DOMSTAL DT (Domperidone dispersible 5 mg and 10 mg tablet) [Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES: Domperidone is a dopamine antagonist and anti-emetic, due to a combination of peripheral (gastrokinetic) effects and antagonism of dopamine receptors in the chemoreceptor trigger zone.

INDICATION: For the relief of post-prandial symptoms of fullness, nausea, epigastric bloating and belching that is occasionally accompanied by epigastric discomfort and heartburn. For the relief of nausea and vomiting of less than 48 hours duration.

DOSAGE AND ADMINISTRATION: The tablet should be dispersed in a teaspoonful of water before administration. For the relief of symptoms of post prandial stomach discomfort: Adults and children 16 years of age and older- Up to 10mg three times daily and at night for maximum duration of 2 weeks. For the relief of nausea and vomiting: Adults and children 16 years of age and older-Up to 10mg three times daily and at night for maximum duration of 48 hours. Use in children below 16 years of age is not recommended.

CONTRAINDICATION: In patients with known hypersensitivity to domperidone or to any component of the formulation. In patients with prolactin-releasing pituitary tumour (prolactinoma), hepatic and/or renal impairment and when stimulation of gastric motility could be harmful (gastrointestinal haemorrhage, mechanical obstruction or perforation).

WARNINGS & PRECAUTIONS: Use is not recommended during lactation, in infants (due to risk of neurological side effects and in patients with hepatic or renal insufficiency. Concomitant use with potent CYP3A4 inhibitors ketoconazole is not recommended due to prolongation of QT interval.

DRUG INTERACTIONS: Interacts with ketoconazole and erythromycin when administered concomitantly.

ADVERSE REACTIONS: Allergic reactions including anaphylaxis, agitation, nervousness, increased prolactin levels, extrapyramidal side effects, convulsion, QT prolongation, ventricular arrhythmias, gastrointestinal disorders, transient intestinal cramps, diarrhea, rash, agitation, somnolence, amenorrhea and liver function test abnormal.

MARKETED BY:



TORRENT PHARMACEUTICALS LTD.

Torrent House, Off Ashram Road,

Ahmedabad-380 009, INDIA

IN/DOMSTAL DT/5,10mg/Mar 2015/01/AbPI

(Additional information is available on request)